

Product datasheet for AP01354PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

CTNND1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Specificity: Catenin-δ1 antibody detects endogenous levels of Catenin-δ1 protein. (region surrounding

Gly224)

Formulation: Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

Purification: Affinity chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 108 kDa

Gene Name: catenin delta 1

Database Link: Entrez Gene 1500 Human

<u>060716</u>




Background:

The α , β , δ , and γ -catenins are proteins that bind to the highly conserved, intracellular cytoplasmic tail of E-cadherin. Together, the catenin/cadherin complexes play an important role mediating cellular adhesion. α -catenin, a 102 kDa protein, interacts with E-cadherin associated protein, and also associates with other members of the cadherin family, such as N-cadherin and P-cadherin. The 92 kDa β -catenin associates with the cytoplasmic portion of E-cadherin, which is necessary for the function of E-cadherin as an adhesion molecule. β -catenin also complexes with the tumor suppressor protein APC. δ -catenin interacts with Presenilin 1 and is expressed in the brain. The gene encoding δ -catenin maps to human chromosome 5p15.2. A hemizygous loss of the gene encoding δ -catenin leads to the mental retardation associated with Cri-du-Chat syndrome. γ -catenin, also known as plakoglobin, is an 80-88 kDa protein that binds α -catenin and N-cadherin. In addition, the transmembrane phosphatase PTPm associates with catenin/cadherin complexes and may regulate complex signaling.


Synonyms:

CTNND1, KIAA0384, p120 catenin, p120(cas), Delta-Catenin

Product images:



Western blot (WB) analysis of Catenin-δ1 antibody (Cat.-No.: AP01354PU-N) in extracts from HuvEc cells.



Immunohistochemistry (IHC) analyzes of Cateninδ1 antibody (Cat.-No.: AP01354PU-N) in paraffinembedded human brain tissue