

## Product datasheet for AP01355PU-N

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **SMAD2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

**Recommended Dilution:** Western Blot: 1/500-1/1000.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

**Specificity:** Smad2 antibody detects endogenous levels of Smad2 protein. (region surrounding Pro246)

Formulation: Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

**Purification:** Affinity chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 52 kDa

Gene Name: SMAD family member 2

Database Link: Entrez Gene 4087 Human

Q15796






#### Background:

Smad proteins, the mammalian homologs of the Drosophila mothers against decapentaplegic (Mad), have been implicated as downstream effectors of TGF\$/BMP signaling. Smad1 (also designated Madr1 or JV4-1) and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF\$\beta\$ and activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to activin/TGF\$\beta\$ signaling by interfering with TGF\$\beta\$-mediated phosphorylation of other Smad proteins.

Synonyms:

SMAD family member 2, SMAD-2, SMAD 2, MADH2, MAD homolog 2, MADR2, Mad-related protein 2, JV18-1

#### **Product images:**



Western blot (WB) analysis of Smad2 antibody (Cat.-No.: AP01355PU-N) in extracts from COLO cells treated with serum 20% 15'