

Product datasheet for AP01356PU-M

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

Syndecan 4 (SDC4) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunofluorescence: 1/50-1/200.

Immunohistochemistry on paraffin sections 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 150-198 of Human Syndecan-4.

Specificity: This antibody detects endogenous levels of Syndecan-4 protein. (region surrounding Lys175)

Formulation: Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 22 kDa

Gene Name: syndecan 4

Database Link: Entrez Gene 6385 Human

P31431





Background:

Syndecans are type I integral membrane proteoglycans that contain both chondroitin sulfate and heparan sulfate groups. Syndecans are involved in cell-extracellular matrix adhesion and growth factor binding. Syndecan-1 (SYND1, also called CD138) is an 85-92 kDa extracellular matrix receptor, which binds to Collagens, Fibronectin and Thrombospondin. Syndecan-1 and Syndecan-3 (also designated N-Syndecan) interact with MK (midkine), a growth/differentiation factor invloved in embryogenesis of the central nervous system. Syndecan-2 (also designated Fibroglycan) is highly expressed at areas of high morphogenetic activity, such as epithelial-mesenchymal interfaces and the prechondrogenic and preosteogenic mesenchymal condensations. Syndecan-4 (also designated Amphiglycan or Ryudocan) functions cooperatively with integrins in the processes of cell spreading, focal adhesion assembly and actin stress fiber assembly.


Synonyms:

SYND4, Amphiglycan, Ryudocan core protein

Product images:



Western blot (WB) analysis of Syndecan-4 antibody in extracts from HepG2 cells.



Immunohistochemistry (IHC) analyzes of Syndecan-4 antibody in paraffin-embedded human brain tissue