

Product datasheet for AP01374PU-N

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

Rockville, MD 20850, US

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CYP7B1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunohistochemistry on Paraffin Sections: 1/50-1/200.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of CYP7B1 protein. (region surrounding Gln127)

Formulation: PBS pH~7.2 with 0.05% Sodium Azide as preservative

State: Aff - Purified

State: Liquid purified Ig fraction (>95% pure by SDS-PAGE)

Concentration: 1.0 mg/ml

Purification: Affinity-Chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 58 kDa

Gene Name: cytochrome P450 family 7 subfamily B member 1

Database Link: Entrez Gene 9420 Human

O75881





Background:

P450 enzymes constitute a family of monooxygenase enzymes that are involved in the metabolism of a wide array of endogenous and xenobiotic compounds including cholesterol. CYP8B1 moderates the ratio of cholic acid over chenodeoxycholic acid to control the solubility of cholesterol.


P450 cholesterol 7-hydroxylase, CYP7A1, is the rate limiting enzyme of bile acid synthesis in the liver, and its expression is mediated by the bile acid receptor FXR. CYP27A1 catalyzes vitamin D3 25-hydroxylation and is localized to the mitochondria in kidney and liver. CYP7B1 (oxysterol 7- α -hydroxylase) functions as an enzyme in the alternate bile acid synthesis pathway. Specifically, CYP7B1 metabolizes 25-and 27-hydroxycholesterol. The gene encoding human CYP7B1 maps to chromosome 8q21.3. Mutations in the CYP7B1 gene may cause a metabolic defect in bile acid synthesis characterized by elevated urinary bile acid excretion, severe cholestasis, cirrhosis and liver synthetic failure.

Synonyms: Cytochrome P450 7B1


Protein Families: Druggable Genome, P450, Transmembrane

Protein Pathways: Primary bile acid biosynthesis

Product images:



Western blot (WB) analysis of CYP7B1 antibody (Cat.-No.: AP01374PU-N) in extracts from LOVO cells.



Immunohistochemistry (IHC) analyzes of CYP7B1 antibody (Cat.-No.: AP01374PU-N) in paraffinembedded human skeletal muscle tissue