

# Product datasheet for AP01386PU-S

# **BMP8A Rabbit Polyclonal Antibody**

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western Blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Immunofluorescence: 1/50-1/200.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	BMP-8A antibody detects endogenous levels of BMP-8A protein. (region surrounding Gln299)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative. State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 45 kDa
Gene Name:	bone morphogenetic protein 8a
Database Link:	<u>Entrez Gene 353500 Human</u> <u>Q7Z5Y6</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** BMP8A Rabbit Polyclonal Antibody – AP01386PU-S


Background:Bone morphogenic proteins (BMPs) are members of the TGFβ superfamily. BMPs are<br/>involved in the induction of cartilage and bone formation. In vivo studies have shown that<br/>BMP-2 (also designated BMP-2A) and BMP-3 can independently induce cartilage formation.<br/>Smad3 association with the TGFβ receptor complex and Smad1 translocation to the nucleus<br/>are observed after the addition of BMP-4 (also designated BMP-2B), suggesting that BMP-4<br/>may play a role in activation of the Smad pathway. BMP-5, BMP-6 and BMP-7 all share high<br/>sequence homology with BMP-2, indicating that they each may be able to induce cartilage<br/>formation. BMP-8 is thought to be involved in early development, as detectable expression<br/>has not been found in adult organs.Two BMP-8 proteins exist, namely BMP-8A and BMP-8B<br/>(also designated OP-2), and are encoded by two distinct genes.

Synonyms: Bone morphogenetic protein 8A, BMP-8A

### **Product images:**



Western blot (WB) analysis of BMP-8A antibody (Cat.-No.: [AP01386PU-N]) in extracts from COLO cells.



Immunohistochemistry (IHC) analyzes of BMP-8A antibody (Cat.-No.: [AP01386PU-N]) in paraffinembedded human brain tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US