

# Product datasheet for AP01387PU-M

## **ARFGEF2** Rabbit Polyclonal Antibody

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western Blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Immunofluorescence: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	BIG2 antibody detects endogenous levels of BIG2 protein. (region surrounding Leu1527)
Formulation:	Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 210 kDa
Gene Name:	ADP ribosylation factor guanine nucleotide exchange factor 2
Database Link:	<u>Entrez Gene 10564 Human</u> <u>Q9Y6D5</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** ARFGEF2 Rabbit Polyclonal Antibody – AP01387PU-M


Background:Guanine nucleotide-exchange proteins (GEPs) accelerate replacement of bound GDP with<br/>GTP and thereby activate ADP-ribosylation factors (ARFs), a family of guanine nucleotide-<br/>binding proteins that play an important role in intracellular vesicular trafficking. GEPs<br/>comprise two major families, large GEPs that are inhibited by brefeldin A (BFA), a protein that<br/>effects golgi structure, and a group of smaller GEPs that are insenstive to BFA. Two genes for<br/>GEPs found on human chromosomes 8 and 20 encode BFA sensitive GEPs of approximately<br/>200 kDa, designated BIG1 and BIG2. Both GEPS contain a sec7 domain that is responsible for<br/>their brefeldin inhibition and also their catalytic activity. In vivo, BIG1 and BIG2 exist in<br/>macromolecular complexes that move between the golgi membranes and cytosol. BIG2<br/>associates with PKA regulatory subunits, implying that BIG2 may act as an A kinase-anchoring<br/>protein (AKAP) that could coordinate the cAMP and ARF regulatory pathways.

Synonyms: ARFGEP2, BIG2, Brefeldin A-inhibited GEP2

#### **Product images:**



Western blot (WB) analysis of BIG2 antibody (Cat.-No.: [AP01387PU-N]) in extracts from A549 cells.



Immunohistochemistry (IHC) analyzes of BIG2 antibody (Cat.-No.: [AP01387PU-N]) in paraffinembedded human colon carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US