

Product datasheet for AP01392PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

CLCN6 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: Western Blot: 1/500-1/1000.

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Specificity: CLC-6 antibody detects endogenous levels of CLC-6 protein. (region surrounding Lys649)

Formulation: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

Purification: Affinity chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 90 kDa

Gene Name: chloride voltage-gated channel 6

Database Link: Entrez Gene 26372 MouseEntrez Gene 1185 Human

P51797






Background:

The family of voltage-dependent chloride channels (CLCs) regulate cellular trafficking of chloride ions, a critical component of all living cells. CLCs regulate excitability in muscle and nerve cells, aid in organic solute transport and maintain cellular volume. The genes encoding human CLC-1 through CLC-7 map to chromosomes 7, 3q26, 4q32, Xp22, Xp11, 1p36 and 16p13, respectively. CLC-1 is highly expressed in skeletal muscle. Mutations in the gene encoding CLC-1 lead to myotonia, an inheritable disorder characterized by muscle stiffness and renal salt wasting. CLC-2 is highly expressed in the epithelia of several organs including lung, which suggests CLC-2 may be a possible therapeutic target for cystic fibrosis. CLC-3 expression is particularly abundant in neuronal tissue, while CLC-4 expression is evident in skeletal and cardiac muscle as well as brain. Mutations in the gene encoding CLC-5 lead to Dent's disease, a renal disorder characterized by proteinuria and hypercalciuria. CLC-6 and CLC-7 are broadly expressed in several tissues including testes, kidney, brain and muscle.

Synonyms: Chloride channel protein 6, CIC-6, CIC6, KIAA0046

Protein Families: Druggable Genome, Ion Channels: Other, Transmembrane

Product images:



Western blot (WB) analysis of CLC-6 antibody (Cat.-No.: AP01392PU-N) in extracts from HT-29