

## **Product datasheet for AP01414PU-M**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Protein C inhibitor (SERPINA5) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: ELISA, IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human
Host: Rabbit

Clonality: Polyclonal

**Specificity:** PAI-3 antibody detects endogenous levels of PAI-3 protein. (region surrounding Leu291)

**Formulation:** Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

**Concentration:** 1.0 mg/ml

**Purification:** Affinity chromatography

Conjugation: Unconjugated

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 42 kDa

Gene Name: serpin family A member 5

Database Link: Entrez Gene 5104 Human

P05154




Background:

PAI-1, PAI-2 and PAI-3 (plasminogen activator inhibitor-1, -2 and -3) are members of the serpin serine proteinase inhibitor family. PAI-1 and PAI-2 regulate uPA (urokinase-type plasminogen activator) and TPA (tissue plasminogen activator), resulting in the inhibition of proteolytic activity. Members of the serpin family generally complex with their target proteinases, then disassociate slowly into cleaved species that fold into stable inactive forms. PAI-1 can fold into the inactive state without cleavage resulting in the latent form of PAI-1. Activity can be restored to the latent form of PAI-1 through denaturation and renaturation. PAI-2 occurs in secreted and cytosolic forms through facultative polypeptide translocation. PAI-3 inhibits plasminogen activators as well as activated protein C. PAI-3 is secreted in plasma, but is also expressed in liver.


Synonyms:

Serpin A5, PAI-3, PAI3, PLANH3, PCI, PROCI

## **Product images:**



Western blot (WB) analysis of PAI-3 antibody (Cat.-No.: [AP01414PU-N]) in extracts from Jurkat cells.



Immunohistochemistry (IHC) analyzes of PAI-3 antibody (Cat.-No.: [AP01414PU-N]) in paraffinembedded human liver carcinoma.