

# Product datasheet for AP01444PU-M

## **IRF3 Rabbit Polyclonal Antibody**

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
21	
Applications:	IHC, WB
Recommended Dilution:	Western Blot: 1/500-1/1000.
	Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	IRF-3 antibody detects endogenous levels of IRF-3 protein. (region surrounding Val391)
Formulation:	Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2. State: Aff - Purified
	State: Liquid purified lg fraction
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 47 kDa
Gene Name:	interferon regulatory factor 3
Database Link:	Entrez Gene 3661 Human
	<u>Q14653</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** IRF3 Rabbit Polyclonal Antibody – AP01444PU-M


Background:Interferon regulatory factor-1 (IRF-1) and IRF-2 have been identified as novel DNA-binding<br/>factors that function as regulators of both type I interferon (interferon-α and β) and<br/>interferon-inducible genes. The two factors are structurally related, particularly in their N-<br/>terminal regions, which confer DNA binding specificity. In addition, both bind to the same<br/>sequence within the promoters of interferon-α and interferon-β genes. IRF-1 functions as an<br/>activator of interferon transcription, while IRF-2 binds to the same ciselements and represses<br/>IRF-1 action. IRF-1 and IRF-2 have been reported to act in a mutually antagonistic manner in<br/>regulating cell growth; overexpression of the repressor IRF-2 leads to cell transformation<br/>while concomitant overexpression of IRF-1 causes reversion. IRF-1 and IRF-2 are members of<br/>a larger family of DNA binding proteins that includes IRF-3, IRF-4, IRF-5, IRF-6, IRF-7, ISGF-3y<br/>p48 and IFN consensus sequence-binding protein (ICSBP).

Synonyms: Interferon regulatory factor 3, IRF-3

### **Product images:**



Western blot (WB) analysis of IRF-3 antibody (Cat.-No.: [AP01444PU-N]) in extracts from HeLa cells.



Immunohistochemistry (IHC) analyzes of IRF-3 antibody (Cat.-No.: [AP01444PU-N]) in paraffinembedded human brain tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US