

Product datasheet for AP01446PU-N

MEK1 (MAP2K1) Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western Blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	MEK1 antibody detects endogenous levels of MEK1 protein. (region surrounding Pro293)
Formulation:	Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 40 kDa
Gene Name:	mitogen-activated protein kinase kinase 1
Database Link:	<u>Entrez Gene 5604 Human</u> <u>Q02750</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Service MEK1 (MAP2K1) Rabbit Polyclonal Antibody – AP01446PU-N


Background:A family of protein kinases located upstream of the MAP kinases and responsible for their
activation has been identified. The prototype member of this family, designated MAP kinase
kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and
tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member,
MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or MKK-3) functions to activate
p38 MAP kinase, and MEK-4 (also called SEK1 or MKK-4) activates both p38 and JNK MAP
kinases. MEK-5 appears to specifically phosphorylate ERK5, whereas MEK-6 phosphorylates
p38 and p38b. MEK-7 (or MKK-7) phosphorylates and activates the JNK signal transduction
pathway.

Synonyms: MAPKK 1, ERK activator kinase 1, MAPK/ERK kinase 1, MEK1, PRKMK1, MAP kinase kinase 1

Product images:



Western blot (WB) analysis of MEK1 antibody (Cat.-No.: AP01446PU-N) in extracts from HUVEC cells.



Immunohistochemistry (IHC) analyzes of MEK1 antibody (Cat.-No.: AP01446PU-N) in paraffinembedded human brain tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US