

# Product datasheet for AP01466PU-N

# **CRABP2** Rabbit Polyclonal Antibody

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western Blot: 1/500-1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of CRABP-II protein. (region surrounding Pro81)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 with 0.05% Sodium Azide as preservative State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~16 kDa
Gene Name:	cellular retinoic acid binding protein 2
Database Link:	<u>Entrez Gene 1382 Human</u> <u>P29373</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** CRABP2 Rabbit Polyclonal Antibody – AP01466PU-N

Background: The cellular retinoic acid-binding protein (CRABP)-I and a related isoform CRABP-II are nuclear receptors for retinoic acid (RA), an important regulator of cell growth and differentiation in fetal and adult tissues. These CRABP proteins mediate the downstream effects of RA in distinct ways. CRABP-I negatively regulates the activity of RA by enhancing the production of RAmetabolizing enzymes and increasing the rate at which RA is degraded. CRABP-II enhances the effects of RA by directly interacting with RA receptors (RAR) and, in turn, promoting the formation of RAR-RA complexes and stimulating RA-mediated gene transcription. Both CRABP-I and CRABP-II are expressed in the embryo, and CRABP-I is ubiquitously expressed in various adult tissues. The expression of CRABP-II is elevated in cells that synthesize relatively large amounts of RA, and it is also predominantly expressed in skin, uterus, ovary, and in the choroid plexus.

Synonyms: CRABP-II

### **Product images:**



Western blot (WB) analysis of CRABP-II antibody (Cat.-No.: AP01466PU-N) in extracts from RAW264.7 cell

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US