

## **Product datasheet for AP01476PU-M**

## OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **GORASP2** Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

Recommended Dilution: Western Blot: 1/500-1/1000.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

Specificity: GRASP55 antibody detects endogenous levels of GRASP55 protein. (region surrounding

Leu215)

**Formulation:** Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

**Concentration:** 1.0 mg/ml

**Purification:** Affinity chromatography

Conjugation: Unconjugated

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 47 kDa

**Gene Name:** golgi reassembly stacking protein 2

Database Link: Entrez Gene 70231 MouseEntrez Gene 113961 RatEntrez Gene 26003 Human

Q9H8Y8





Background:

The Golgi apparatus is a highly complex organelle comprised of a stack of cisternal membranes on the secretory pathway from the ER to the cell surface. The structure is maintained by an exoskeleton or Golgi matrix constructed from a family of coiled-coil protein, the golgins and other peripheral membrane components such as GRASP55 and GRASP65. GRASP55 (Golgi reassembly stacking protien or p59) is a component of the Golgi stacking machinery. GRASP55 is highly homologous to GRASP65 and contains two PDZ domains. GRASP55 is myristoylated and palmitoylated. Unlike GRASP65, GRASP55 does not have detectable binding with the vesicle docking protein GM130 and is located on the medial-Golgi rather than cis-Golgi. Both GRASP55 and GRASP65 function in the stacking of Golgi cisternae. The novel coiled-coil protein golgin 45 interacts with GRASP55 and the GTP form of Rab 2, suggesting that GRASP55 and golgin 45 form a Rab 2 effector complex on medial-Golgi essential for normal protein transport and Golgi structure. ERK2 directly phosphorylates GRASP55, which is phosphorylated in mitotic cells, suggesting that mitogen-activated protein kinase kinase (MKK)/ERK pathway phosphorylates the Golgi during mitosis.


Synonyms:

Golgi reassembly-stacking protein 2, GRS2, Golgi phosphoprotein 6, p59, Golgi marker

**Protein Families:** 

Druggable Genome

## **Product images:**



Western blot (WB) analysis of GRASP55 antibody (Cat.-No.: [AP01476PU-N]) in extracts from LOVO cells.