

Product datasheet for AP01479PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GNL1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: Western Blot: 1/500-1/1000.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Specificity: GNL1 antibody detects endogenous levels of GNL1 protein. (region surrounding Gln103)

Formulation: Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

Purification: Affinity chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 69 kDa

Gene Name: G protein nucleolar 1 (putative)

Database Link: Entrez Gene 2794 Human

P36915






Background:

GNL1 (guanine nucleotide-binding protein-like 1) is a nuclear protein that likely acts as a regulator of the histocompatibility cluster. GNL1 and MMR1, the murine homolog, are localized within or close to the MHC class I region and belong to the MMR1/HSR1 GTP-binding protein family. GTPases, such as GNL1, from the MMR1/HSR1 GTP-binding protein subfamily are circularly rearranged G-motifs that play a critical role in maintaining normal cell growth. Deletion of these genes results in severe growth defects with a marked reduction in mature rRNA species and a concomitant accumulation of the 35S pre-rRNA transcript. Deletion also causes the ribosomal protein RPL25A to fail exportation from the nucleolus. Deletion of any of the G-domain motifs will result in a null phenotype and nuclear/nucleolar localization that lacks the nucleolar export of preribosomes and is accompanied by a distortion of the nucleolar structure. Upon DNA damage GNL1 is phosphorylated by a kinase, possibly Atm or ATR.

Synonyms:

Guanine nucleotide-binding protein-like 1, GTP-binding protein HSR1

Product images:



Western blot (WB) analysis of GNL1 antibody (Cat.-No.: [AP01479PU-N]) in extracts from K562 cells.