

Product datasheet for **AP01497PU-M**

CHURC1-FNTB Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western Blot: 1/500-1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	FT β antibody detects endogenous levels of FT β protein. (region surrounding Glu285)
Formulation:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 50 kDa
Gene Name:	CHURC1-FNTB readthrough
Database Link:	Entrez Gene 100529261 Human P49356




[View online »](#)

Background:

Mammalian protein farnesyl transferases are heterodimeric proteins containing two nonidentical α and β subunits that attach farnesyl residues to a cysteine at the fourth position from the COOH terminus of several proteins, including nuclear lamins and p21Ras proteins. The natural substrates contain the Cys-A-A-Xaa recognition sequence, where the A residues are aliphatic and Xaa represents methionine, serine, glutamine or cysteine. The purified farnesyl transferase is an α - β heterodimer. The β subunit binds the peptide substrate while the α subunit is suspected to participate in formation of a stable complex with the substrate farnesyl pyrophosphate. The α subunit is shared with a second prenyl transferase, geranyl-geranyl transferase, that attaches 20 carbon geranylgeranyl to Ras related proteins that terminate in a Cys-A-A-Xaa recognition site in which Xaa is leucine.

Synonyms:

FTase-beta, CAAX farnesyltransferase subunit beta

Product images:


Western blot (WB) analysis of FT β antibody (Cat.-No.: [AP01497PU-N]) in extracts from RAW264.7 cells.