

# Product datasheet for AP01505PU-N

# **CACNA1H Rabbit Polyclonal Antibody**

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western blot: 1/500-1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 461-510 of Human T-type Ca++ CP $lpha$ 1H.
Specificity:	This antibody detects endogenous levels of T-type C $a^+$ CP $\alpha$ 1H protein. (region surrounding Pro492)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified lg fraction (> 95% pure by SDS-PAGE) Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~259 kDa
Gene Name:	calcium voltage-gated channel subunit alpha1 H
Database Link:	<u>Entrez Gene 8912 Human</u> <u>O95180</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** CACNA1H Rabbit Polyclonal Antibody – AP01505PU-N

Background:Voltage-dependent Ca++ channels mediate Ca++ entry into excitable cells in response to<br/>membrane depolarization, and they are involved in a variety of Ca++-dependent processes,<br/>including muscle contraction, hormone or neurotransmitter release and gene expression.<br/>Calcium channels are highly diverse, multimeric complexes composed of an α1 subunit, an<br/>intracellular β subunit, a disulfide linked α2/δ subunit and a transmembrane y subunit. Ca++<br/>currents are characterized on the basis of their biophysical and pharmacologic properties<br/>and include L-, N-, T-, P-, Q-, and R- types. T-type Ca++ currents are activated and inactivated<br/>more rapidly and at more negative membrane potentials than other Ca++ current types. T-<br/>type Ca++ channels enhance odor sensitivity by lowering the threshold of spike generation in<br/>olfactory receptor cells (ORCs).

Synonyms: Cav3.2

### **Product images:**



Western blot (WB) analysis of T-type Ca++ CP aH antibody in extracts from A549 cells.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., <u>9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US</u>