

Product datasheet for AP01506PU-N

ADCY4 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	Western Blot: 1/500-1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	A Cyclase IV antibody detects endogenous levels of A Cyclase IV protein. (region surrounding Leu226)
Formulation:	Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 120 kDa
Gene Name:	adenylate cyclase 4
Database Link:	<u>Entrez Gene 196883 Human</u> <u>Q8NFM4</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE ADCY4 Rabbit Polyclonal Antibody – AP01506PU-N

Background:Adenylyl cyclases function to convert ATP to cyclic AMP in response to activation by a variety
of hormones, neurotransmitters and other regulatory molecules. Cyclic AMP, in turn, activates
several other target molecules to control a broad range of diverse phenomena such as
metabolism, gene transcription and memory. Adenylyl cyclases respond to receptor-initiated
signals, mediated by the Gs and Gi heterotrimeric G proteins. The binding of an agonist to a
Gs-coupled receptor catalyzes the exchange of GDP (bound to Gα s) for GTP, the dissociation
of GTP-Gα s from Gβγ and Gα s-mediated activation of adenylyl cyclase. Adenylyl cyclase IV
(AC IV) and IX mRNA are expressed in all kidney nephron segments. AC IV exhibits moderate
staining in type II and type IV fibrocytes in rat cochlea and immunoreactivity is also observed
in type I fibrocytes. Activation of the D2 dopaminergic and m4 muscarine receptors inhibits
the activity of adenylyl cyclase isozymes I, V, VI and VIII, whereas type II, IV and VII are
stimulated and type III is not affected.

Synonyms:

Adenylate cyclase type IV, Adenylyl cyclase 4, ADCY4

Product images:



Western blot (WB) analysis of A Cyclase IV antibody (Cat.-No.: AP01506PU-N) in extracts from Jurkat cells.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US