

# Product datasheet for AP01532PU-M

# ATF2 pThr73/55 Rabbit Polyclonal Antibody

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, IP, WB
Recommended Dilution:	Western Blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Immunoprecipitation: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	p-ATF2 (pThr73/55) antibody detects endogenous levels of p-ATF2 protein.
Formulation:	Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	activating transcription factor 2
Database Link:	<u>Entrez Gene 1386 Human</u> <u>P15336</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### CRIGENE ATF2 pThr73/55 Rabbit Polyclonal Antibody – AP01532PU-M


Background:Eukaryotic gene transcription is regulated by sequence-specific transcription factors which<br/>bind modular CIS acting promotor and enhancer elements. The cAMP response element<br/>(CRE), one of the best studied of such elements, consists of the palindromic octanucleotide<br/>TGACGTCA. Several CRE binding proteins have been identified within the ATF/CREB family, the<br/>best characterized of which include CREB-1, CREB-2 (also designated ATF-4), ATF-1, ATF-2 and<br/>ATF-3. These proteins share highly related COOH terminal leucine zipper dimerization and<br/>basic DNA binding domains but are highly divergent in their amino terminal domains.<br/>Although each of the ATF/CREB proteins appear capable of binding CRE in its homodimeric<br/>form, certain of these also bind as heterodimers, both within the ATF/CREB family and even<br/>with members of the AP-1 transcription factor family.

Synonyms: ATF-2, CREB2, CREBP1, Activating transcription factor 2, CRE-BP1, HB16, CREB-2

## **Product images:**



Western blot (WB) analysis of p-ATF2 (pThr73/55) antibody (Cat.-No.: [AP01532PU-N]) in extracts from 293 cells treated with UV 15'.



Immunohistochemistry (IHC) analyzes of p-ATF2 (pThr73/55) antibody (Cat.-No.: [AP01532PU-N]) in paraffin-embedded human breast carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US