

## Product datasheet for AP01596PU-M

## OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## GSK3 beta (GSK3B) pSer9 Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, IP, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunoprecipitation: 1/50-1/200.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of GSK3b pSer9 protein.

**Formulation:** Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Concentration: 1.0 mg/ml

**Purification:** Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: One year from despatch.

**Predicted Protein Size:** ~ 48 kDa

**Gene Name:** glycogen synthase kinase 3 beta

**Database Link:** Entrez Gene 2932 Human

P49841




Background:

Glycogen synthase kinase 3, or GSK-3, is a serine/threonine, proline-directed kinase involved in a diverse array of signaling pathways, including glycogen synthesis and cellular adhesion, and has been implicated in Alzheimer's disease. Two forms of GSK-3, designated GSK-3 $\alpha$  and GSK-3 $\beta$ , have been identified and differ in their subcellular localization. Tau, a microtubule-binding protein which serves to stabilize microtubules in growing axons, is found to be hyper-phosphorylated in paired helical filaments (PHF), the major fibrous component of neurofibrillary lesions associated with Alzheimer's disease. Hyperphosphorylation of Tau is thought to be the critical event leading to the assembly of PHF. Six Tau protein isoforms have been identified, all of which are phosphorylated by GSK-3. This presents the possibility that miscues in GSK-3 signaling contribute to the onset of Alzheimer's disease.


Synonyms:

GSK-3 beta, Glycogen synthase kinase-3 beta, GSK3B

## **Product images:**



Western blot (WB) analysis of GSK3 $\beta$  pSer9 antibody (Cat.-No.: [AP01596PU-N]) in extracts from HeLa cells treated with EGF.



Immunohistochemistry (IHC) analysis of GSK3b pSer9 antibody (Cat.-No.: [AP01596PU-N]) in paraffin-embedded human breast carcinoma tissue.