

## **Product datasheet for AP01598PU-M**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **HDAC5** pSer498 Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

**Reactivity:** Human, Mouse

**Host:** Rabbit

Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of HDAC5 pSer498 protein.

**Formulation:** Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

**Concentration:** 1.0 mg/ml

**Purification:** Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: One year from despatch.

Predicted Protein Size: ~ 122 kDa

**Gene Name:** histone deacetylase 5

**Database Link:** Entrez Gene 10014 Human

Q9UQL6




Background:

HDAC5 is a member of the class II mammalian histone deacetylase family, which is structurally related to yeast HDA1. Human HDAC5 is composed of 1122 amino acid residues. The deacetylase domain of HDAC5 is located at the C-terminal half of the molecule. The N-terminal non-deacetylase domain does not show any significant homology with any published sequence. Both domains are required for HDAC5-mediated repression of gene transcription. HDAC5 interacts with a growing number of transcriptional factors including MEF2A as well as other HDAC proteins. The interacting complexes bind to specific regions of chromatin and regulate gene transcription in these regions.


Synonyms:

Histone deacetylase 5, HD5, KIAA0600, Antigen NY-CO-9

## **Product images:**



Western blot (WB) analysis of HDAC5 pSer498 antibody (Cat.-No.: [AP01598PU-N]) in extracts from NIH/3T3 cells.



Immunohistochemistry (IHC) analysis of HDAC5 pSer498 antibody (Cat.-No.: [AP01598PU-N]) in paraffin-embedded human breast carcinoma tissue.