

# **Product datasheet for AP01606PU-S**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

#### H3FA (HIST1H3A) pSer10 Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

**Recommended Dilution:** Western Blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Specificity:** This antibody detects endogenous levels of Histone H3.1 pSer10 protein.

**Formulation:** Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

**Concentration:** 1.0 mg/ml

**Purification:** Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: One year from despatch.

**Predicted Protein Size:** ~ 15 kDa

Gene Name: histone cluster 1, H3a

Database Link: Entrez Gene 8350 Human

P68431



Synonyms:

#### H3FA (HIST1H3A) pSer10 Rabbit Polyclonal Antibody - AP01606PU-S

**Background:** Histones are basic nuclear proteins that together with DNA make up the nucleosome


structure in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin structure is further compacted through the interaction of a linker histone H1. Methylation of specific residues in the histone tails is a central modification for regulating epigenetic transitions in chromatin. Modified lysine residues can exist in a mono-,

di-, or tri-methylated state, whilst arginine residues can be mono- or di-methylate


H3/a, H3/b, H3/c, H3/d, H3/f, H3/h, H3/i, H3/j, H3/k, H3/l, HIST1H3A, H3FA, HIST1H3B,

HIST1H3C, HIST1H3D, HIST1H3E, HIST1H3F, HIST1H3G, HIST1H3H, HIST1H3I, HIST1H3J

### **Product images:**



Western blot (WB) analysis of Histone H3.1 pSer10 antibody (Cat.-No.: [AP01606PU-N]) in extracts from HeLa cells.



Immunohistochemistry (IHC) analyzes of Histone H3.1 pSer10 antibody (Cat.-No.: [AP01606PU-N]) in paraffin-embedded human skeleta I muscle tissue.