

Product datasheet for AP01629PU-N

LIM kinase 2 (LIMK2) pThr505 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: IF, IHC, WB Recommended Dilution: Western Blot: 1/500-1/1000. Immunohistochemistry on paraffin sections 1/50-1/200. Immunofluorescence: 1/50-1/200. **Reactivity:** Human, Mouse, Rat Host: Rabbit **Clonality:** Polyclonal Specificity: This antibody detects endogenous levels of LIMK2 pThr505 protein. Formulation: Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative. State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). **Concentration:** 1.0 mg/ml **Purification:** Affinity Chromatography using epitope-specific immunogen. **Conjugation:** Unconjugated Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage: Avoid repeated freezing and thawing. Shelf life: One year from despatch. Stability: Predicted Protein Size: ~ 72 kDa Gene Name: LIM domain kinase 2 Database Link: Entrez Gene 3985 Human P53671



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


	LIM kinase 2 (LIMK2) pThr505 Rabbit Polyclonal Antibody – AP01629PU-N
Background:	Proteins containing LIM motifs are typically involved in cell fate determination and growth control. A family of proteins designated LIM kinases, incuding LIMK-1 and LIMK-2, has been identified. LIMK-1 has been shown to regulate the stabilization of F-Actin structures and Cofilin activity, indicating that LIMK-1 plays a role in a signaling pathway involved in the regulation of cell motility and morpho-genesis. LIMK-1 inhibits neuronal differentiation of PC12 cells, and it is thought to act by interfering with events downstream of MAPK activation. Expression patterns of LIMK-1 and LIMK-2 suggest that these proteins may have different functions during development. A truncated form of LIMK-2 has been identified in adult testis that is thought to arise from an alternative initiation exon.

Synonyms: LIMK2, LIM domain kinase 2, LIMK-2, EC=2.7.11.1

Product images:



Western blot (WB) analysis of LIMK2 pThr505 antibody (Cat.-No.: AP01629PU-N) in extracts from COLO cells.



Immunohistochemistry (IHC) analysis of LIMK2 pThr505 antibody (Cat.-No.: AP01629PU-N) in paraffin-embedded human breast carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US