

Product datasheet for AP01632PU-N

MEF2A pThr312 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IHC, IP, WB

Recommended Dilution: ELISA: 1/20000-1/40000.

Western Blot: 1/500-1/1000.

Immunohistochemistry: 1/50-1/200. Immunoprecipitation: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of MEF2A protein only when phosphorylated at

Thr312.

Formulation: Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: myocyte enhancer factor 2A

Database Link: Entrez Gene 4205 Human

002078



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

MEF-2 is a muscle-specific DNA binding protein that recognizes an A+T-rich sequence [CTA (A/T)4 TAG] localized in the control regions of numerous muscle-specific genes. MEF-2 belongs to the MADS (MCM1, Agamous, Deficiens and serum-response factor) box family of transcription factors. The MEF-2 proteins arise from several alternatively spliced isoforms of the MEF-2 gene. MEF-2 expression is ubiquitous, but appears to be preferential in skeletal and cardiac muscle cells. Phosphorylation of different MEF-2C isoforms affects their expression pattern and transactivation function. Big MAP kinase 1 (BMK1) enhances the transactivation activity of MEF-2C by phosphorylating Ser 387. Serum is a potent stimulator of BMK1-induced MEF-2C phosphorylation. p38 MAPK can phosphorylate MEF-2C at positions Ser 387, Thr 293 and Thr 300. Phosphorylation of MEF-2C by either p38 MAPK or ERK5/BMK1 is necessary for Smad-MEF-2 signaling cooperativity.


Synonyms: MEF2, Myocyte-specific enhancer factor 2A

Note: Molecular Weight: ~55.0 kDa

Product images:



Western blot (WB) analysis of MEF2A pThr312 antibody in extracts from NIH/3T3 cells.



Immunohistochemistry (IHC) analysis of MEF2A pThr312 antibody in paraffin-embedded human breast carcinoma tissue.