

## Product datasheet for AP01646PU-N

## NFKB1 pSer907 Rabbit Polyclonal Antibody

**Product data:** 

**Product Type: Primary Antibodies** 

IHC, IP, WB **Applications:** 

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunoprecipitation: 1/50-1/200.

Reactivity: Human Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of NFkB-p105/p50 pSer907 protein.

Formulation: Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Concentration: 1.0 mg/ml

**Purification:** Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Shelf life: One year from despatch. Stability:

**Predicted Protein Size:** ~ 110 kDa

Gene Name: nuclear factor kappa B subunit 1

**Database Link:** Entrez Gene 4790 Human

P19838



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Proteins encoded by the v-Rel viral oncogene and its cellular homolog, c-Rel, are members of a family of transcription factors that include the two subunits of the transcription factor NF $\kappa$ B (p50 and p65) and the Drosophila maternal morphogen, dorsal. Both proteins specifically bind to DNA sequences that are the same or slight variations of the 10 bp  $\kappa$ B sequence in the immunoglobulin  $\kappa$  light chain enhancer. This same sequence is also present in a number of other cellular and viral enhancers. The DNA binding activity of NF $\kappa$ B is activated and NF $\kappa$ B is subsequently transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct proteins of the same size have been described, designated p105 and p100. The p105 precursor contains p50 at its N-terminus and a C-terminal region that when expressed as a separate molecule, designated pdI, binds to p50 and regulates its activity


Synonyms:

NFKB1, KBF1, EBP-1, EBP1, NF-kappa-B p50

## **Product images:**



Western blot (WB) analysis of NF?B-p105/p50 pSer907 antibody (Cat.-No.: AP01646PU-N) in extracts from HeLa cells treated with TNF-alpha.



Immunohistochemistry (IHC) analysis of NF?B-p105/p50 pSer907 antibody (Cat.-No.: AP01646PU-N) in paraffin-embedded human breast carcinoma tissue.