

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for AP01659PU-M

### p53 (TP53) pSer46 Rabbit Polyclonal Antibody

#### **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western Blot: 1/500-1/1000.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	p53 antibody detects endogenous levels of p53 pSer46 protein.
Formulation:	Phosphate buffered saline (PBS), pH~7.2 containing 15 mM Sodium Azide as preservative. State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Predicted Protein Size:	~ 46 kDa
Gene Name:	tumor protein p53
Database Link:	<u>Entrez Gene 7157 Human</u> <u>P04637</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ENE p	53 (TP53) pSer46 Rabbit Polyclonal Antibody – AP01659PU-M
und:	p53, a DNA-binding, oligomerization domain and transcription activation domain-containing tumor suppressor, upregulates growth arrest and apoptosisrelated genes in response to stress signals, thereby influencing programmed cell death, cell differentiation and cell cycle control mechanisms. p53 localizes to the nucleus, yet can be chaperoned to the cytoplasm by the negative regulator MDM2, an E3 ubiquitin ligase that is upregulated in the presence of active p53, where MDM2 poly-ubiquitinates p53 for proteasome targeting. p53 fluctuates between latent and active (DNA-binding) conformations and is differentially activated throug posttranslational modifications including phosphorylation and acetylation. Mutations in the DNA-binding domain (DBD, amino acids 110-286) of p53 can compromise energetically favorable association with cis elements and are implicated in several human cancers.
	stress signals, thereby influencing programmed cell death, cell differenti control mechanisms. p53 localizes to the nucleus, yet can be chaperoned the negative regulator MDM2, an E3 ubiquitin ligase that is upregulated active p53, where MDM2 poly-ubiquitinates p53 for proteasome targetin between latent and active (DNA-binding) conformations and is differenti posttranslational modifications including phosphorylation and acetylatic DNA-binding domain (DBD, amino acids 110-286) of p53 can compromis

Synonyms:

Cellular tumor antigen p53, Tumor suppressor p53, Phosphoprotein p53, NY-CO-13

## **Product images:**



Hela whole cell lysate p-p53 (S46) pAb at 1:500 dilution Western blot (WB) analysis of p-p53 (pSer46) antibody Cat.-No.: [AP01659PU-N] in extracts from hela cells.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US