

Product datasheet for AP01710PU-M

oduct datasileet for APO17 for O-IVI

MAPT / TAU pThr205 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of Tau pThr205 protein.

Formulation: Phosphate buffered saline (PBS), pH~7.2 containing 0.05% Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Predicted Protein Size: ~ 78 kDa

Gene Name: microtubule associated protein tau

Database Link: Entrez Gene 4137 Human

P10636



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

MAPT / TAU pThr205 Rabbit Polyclonal Antibody – AP01710PU-M


Background:

Tau is a neuronal microtubule associated protein found predominantly on axons. The function of Tau is to promote tubulin polymerisation and stabilise microtubules, but it also serves to link certain signalling pathways to the cytoskeleton. Tau, in its hyperphosphorylated form, is the major component of paired helical filaments (PHF) and neurofibrillary lesions in Alzheimer's disease (AD) brain. Hyperphosphorylation impairs the microtubule binding function of Tau, resulting in the destabilisation of microtubules in AD brains, ultimately leading to the degeneration of the affected neurons. Hyperphosphorylated tau is also found in a range of other central nervous system disorders. Numerous serine/threonine kinases, including GSK3 beta, PKA, Cdk5, and casein kinase II can phosphorylate Tau.


Synonyms:

MAPTL, MTBT1, Microtubule-associated protein tau, PHF-tau, Neurofibrillary tangle protein, Paired helical filament-tau

Product images:



Western blot (WB) analysis of Tau pThr205 antibody (Cat.-No.: [AP01710PU-N]) in extracts from mouse brain.



Immunohistochemistry (IHC) analyzes of Tau pThr205 antibody (Cat.-No.: [AP01710PU-N]) in paraffin-embedded human breast carcinoma tissue.