

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for AP01745PU-M

### HDAC4 pSer632 Rabbit Polyclonal Antibody

#### **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western Blot: 1/500-1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of HDAC4 pSer632 protein.
Formulation:	Phosphate buffered saline (PBS), pH~7.2 containing 15 mM Sodium Azide as preservative. State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Predicted Protein Size:	~ 119 kDa
Gene Name:	histone deacetylase 4
Database Link:	<u>Entrez Gene 9759 Human</u> <u>P56524</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ORIGENE HDA	C4 pSer632 Rabbit Polyclonal Antibody – AP01745PU-M
ckground:	In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation, and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino-terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, p300/CBP, PCAF (p300/CBPassociated factor), HAT1 and the TFIID subunit TAF II p250. Mammalian HDAC1 (also designated HD1), HDAC2 (also designated RPD3) and HDAC3-6 have been identified as histone deacetylases.
	transcriptional regulation, and a major source of this remodeling is brought about acetylation of nucleosomal histones. Acetylation of lysine residues in the amino-ter domain of histone results in an allosteric change in the nucleosomal conformation increased accessibility to transcription factors by DNA. Conversely, the deacetylatio histones is associated with transcriptional silencing. Several mammalian proteins h identified as nuclear histone acetylases, including GCN5, p300/CBP, PCAF (p300/CBPassociated factor), HAT1 and the TFIID subunit TAF II p250. Mammalian F

Synonyms: Histone deacetylase 4, HD4, KIAA0288

## **Product images:**



Western blot (WB) analysis of p-HDAC4 (pSer632) antibody (Cat.-No.: [AP01745PU-N]) in extracts from Jurkat cells.

Jurkat whole cell lysate p-HDAC4 (S632) pAb at 1:500 dilution

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US