

## Product datasheet for **AP02074SU-N**

### CCL14 (8-15) Rabbit Polyclonal Antibody

#### Product data:

|                        |  |
|------------------------|--|
| Product Type:          | Primary Antibodies   |
| Applications:          | ELISA  |
| Recommended Dilution:  | ELISA (1:1,000).   |
| Reactivity:            | Human  |
| Host:                  | Rabbit   |
| Clonality:             | Polyclonal   |
| Immunogen:             | Synthetic human Chemokine CC-1/3 (aa 8-15) poly Lysinconjugated  |
| Specificity:           | This antibody detects Synthetic human Chemokine CC-1/3 (aa 8-15).  |
| Formulation:           | State: Serum<br>State: Lyophilized serum   |
| Reconstitution Method: | Restore in aqua bidest to initial volume.  |
| Conjugation:           | Unconjugated   |
| Storage:               | Store lyophilized at 2 - 8 °C and reconstituted at -20 °C. Avoid repeated freezing and thawing.  |
| Stability:             | Shelf life: One year from despatch.  |
| Gene Name:             | C-C motif chemokine ligand 14  |
| Database Link:         | <a href="#">Entrez Gene 6358 Human Q16627</a>  |
| Background:            | The CC cytokines are secreted proteins characterized by two adjacent cysteines. The cytokine encoded by this gene induces changes in intracellular calcium concentration and enzyme release in monocytes. This gene expresses both monocistronic and bicistronic transcripts. Bicistronic transcripts include the upstream cytokine gene CCL15. In addition, CCL14 undergoes alternative splicing in the bicistronic transcripts; it is unknown if its monocistronic transcript is also subject to alternative splicing. |
| Synonyms:              | C-C motif chemokine 14, Small-inducible cytokine A14, Chemokine CC-1/CC-3, HCC-1/HCC-3, NCC-2, SCYA14  |
| Note:                  | LocusID 6358   |



[View online »](#)