

# Product datasheet for AP02094SU-N

### Transglutaminase 3 (TGM3) Rabbit Polyclonal Antibody

### **Product data:**

Product Type:	Primary Antibodies
Applications:	ELISA, IF, IHC
Recommended Dilution:	ELISA (Ref.1) Immunofluorescence. Immunohistochemistry on Cryosections: 1/100-1/3200 (Ref.1,2).
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Recombinant Human epidermal Transglutaminase [TGe]
Specificity:	This antibody detects Epidermal Transglutaminase [TGe].
Formulation:	State: Serum State: Lyophilized Serum
<b>Reconstitution Method:</b>	Restore in aqua bidest to initial volume.
Conjugation:	Unconjugated
Storage:	Store lyophilized at 2-8°C and reconstituted at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	transglutaminase 3
Database Link:	<u>Entrez Gene 7053 Human</u> <u>Q08188</u>
Background:	Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved, they all have the same amino acid sequence at their active sites and their activity is calcium-dependent. The protein encoded by this gene consists of two polypeptide chains activated from a single precursor protein by proteolysis. The encoded protein is involved the later stages of cell envelope formation in the epidermis and hair follicle. [provided by RefSeq]
Synonyms:	TGase, TGE
Note:	LocusID 7053



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


Α

Protein Families:

Druggable Genome

## **Product images:**







Immunofluorescence of TGe staining in Cryosection of Human skin of dermatitis herpetiformis patient. The section was incubated with AP02094SU (1/100) followed by incubation with Cy3-labeled Goat ant-Rabbit antibody. This antibody stains the epidermis and aggregates within the dermal papillae (Red). TGe colocalizes with IgA (Green) in these precipitates. Sardy M et al. (2002) J Exp Med 195 (6): 747-757.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US