

## Product datasheet for AP02233SU-S

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Thymosin beta 4 (TMSB4X) (1-14) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, IHC

Recommended Dilution: ELISA: 1/4000 (Ref.1,2).

**Dot Blot:** 1/500.

Immunohistochemistry on Paraffin Sections: 1/200 (Ref.3-7). Immunohistochemistry on Vibratome Sections: 1/100 (Ref.3-7). Immunohistochemistry on Cryo Sections: 1/8000 (Ref.3-7). Immunohistochemistry on Whole-mount: 1/100 (Ref.3-7).

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic Human Thymosin beta-4 (aa 1-14) KLH conjugated

**Specificity:** This antibody detects Thymosin beta-4 (aa 1-14).

There was no cross reactivity obtained with Human Thymosin α1 and Human Thymulin, Tb4

(aa 1-4), Tb9, Tb9 (aa 1-14), AcADKP, ADKP, AcGDKP, AcTDKP, AcSPKD, AcVDKP, BSA.

Formulation: State: Serum

State: Lyophilized Serum

**Reconstitution Method:** Restore in aqua bidest to initial volume.

**Conjugation:** Unconjugated

**Storage:** Store lyophilized at 2-8°C for 6 months or at -20°C long term.

After reconstitution store the antibody undiluted at 2-8°C for one month

or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: thymosin beta 4, X-linked

Database Link: Entrez Gene 7114 Human

P62328





**Background:** Thymosin beta-4 was shown to be essential for all aspects of coronary vessel development in

mice; it stimulates significant outgrowth from quiescent adult epicardial explants, restoring


pluripotency and triggering differentiation of fibroblasts, smooth muscle cells, and

endothelial cells.


Synonyms: T beta-4, TMSB4X, TB4X, THYB4, TMSB4, Seraspenide

Note: LocusID 7114

## **Product images:**



ELISA: Standard curve showing measurement of Thymosin beta-4 in a competitive Immunoassay using AP02233SU. OD was deteremined at 450 nm. The shown standard curve is for demonstration purposes only and can be influenced by Buffer, incubation conditions or dilution of the conjugate.



Immunohistochemistry using AP02233SU Thymosin beta-4 Antibody: Staining of Paraffin Sections of breast cancer tissue. Sections were incubated with AP02233SU (1/200) for 2 h followed by incubation with Aexa488 Goat anti rabbit antibody (1/400). The section were stained for DNA using bis-Benzimide Image A: AP02233SU stains scattered tumor cells strongly. Image B: Antiserum preabsorbed with the antigen does not stain the tumor tissue. Larsson LI et al. (2007) Hum pathol 38 (1):114-9.