

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AP02390PU-N

p53 (TP53) pSer15 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500 - 1/1000; Incubate membrane with diluted antibody in 5% nonfat milk, 1X TBS, 0,1% Tween-20 at 4°C with gentle shaking, overnight. Immunohistochemistry on paraffin sections: 1/50 - 1/100. Immunofluorescence: 1/100 - 1/200.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic phosphopeptide derived from human p53 around the phosphorylation site of serine 15 (P-L-SP-Q-E).
Specificity:	This antibody detects endogenous levels of p53 only when phosphorylated at Serine 15.
Formulation:	PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol as stabilizer State: Aff - Purified State: Liquid purified lg fraction
Concentration:	lot specific
Purification:	Immunoaffinity chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody (in aliquots) at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	tumor protein p53
Database Link:	<u>Entrez Gene 7157 Human</u> <u>P04637</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE p53 (TP53) pSer15 Rabbit Polyclonal Antibody – AP02390PU-N

Background:Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis
depending on the physiological circumstances and cell type. Involved in cell cycle regulation
as a trans-activator that acts to negatively regulate cell division by controlling a set of genes
required for this process. One of the activated genes is an inhibitor of cyclin-dependent
kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS
antigen expression, or by repression of Bcl-2 expression. Implicated in Notch signaling cross-
over. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-
inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth
suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis.

Synonyms:

Cellular tumor antigen p53, Tumor suppressor p53, Phosphoprotein p53, NY-CO-13

Product images:




Figure 2. Western blot analysis of the extracts from HeLa cells untreated or treated with hydroxyurea, using p53 antibody (Line 1 and 2) and p53 (phospho-Ser15) antibody (Line 3 and 4).





Figure 1. Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using p53 (phospho- Ser15) antibody.

P-Peptide

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Immunofluorescence staining of methanol-fixed HeLa cells using p53 (pSer15) Antibody

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US