

## Product datasheet for AP02408PU-S

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **CDC25C Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

Recommended Dilution: Immunohistochemistry on paraffin sections: 1/50 - 1/100.

Western Blot: 1/500 - 1/1000; Incubate membrane with diluted antibody in 5% nonfat milk,

1X TBS, 0,1% Tween-20 at 4°C with gentle shaking, overnight.

Immunofluorescence: 1/100 - 1/200.

Reactivity: Human
Host: Rabbit

Clonality: Polyclonal

**Immunogen:** Synthetic phosphopeptide derived from human cdc25C around the phosphorylation site of

Serine 216 (S-P-S*p*-M-P).

**Specificity:** This antibody detects endogenous levels of CDC25C only when phosphorylated at Serine 216.

Formulation: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol

State: Aff - Purified

State: Liquid purified Ig fraction

**Concentration:** lot specific

**Purification:** Immunoaffinity Chromatography

**Conjugation:** Unconjugated

Storage: Store the antibody (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: One year from despatch.

**Gene Name:** cell division cycle 25C

Database Link: Entrez Gene 995 Human

P30307






Background:


Cdc25C is a tyrosine phosphatase and belongs to the Cdc25 phosphatase family. It has been highly conserved during evolution and it plays a key role in the regulation of cell division. It directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It is also thought to suppress p53-induced growth arrest. Cdc25C is mainly expressed in G2 phase. Multiple alternatively spliced transcript variants of this gene have been described, however, the full-length nature of many of them is not known.

Synonyms: CDC25

## **Product images:**




Western Blot analysis of extracts from 293 cells untreated or treated with serum using cdc25c (pSer216) antibody



Immunohistochemical staining of paraffinembedded human breast carcinoma tissue using cdc25c (pSer216) antibody (left) or the same antibody preincubated with blocking peptide (right)





Immunofluorescence staining of methanol-fixed HeLa cells using cdc25c (pSer216) antibody