

Product datasheet for **AP02463PU-N**

Amyloid Precursor Protein (APP) pThr668 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Suitable for use in Suitable for use in Western blot (1/500-1/1000).
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The antiserum was produced against synthesized phosphopeptide derived from human APP around the phosphorylation site of threonine 668 (A-V-T _p -P-E).
Specificity:	The antibody detects endogenous levels of APP only when phosphorylated at Threonine 668.
Formulation:	PBS (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol. State: Aff - Purified State: Liquid purified Ig fraction.
Concentration:	lot specific
Purification:	Immunoaffinity Chromatography using epitope-specific phosphopeptide. Non-phosphopeptide antibodies was removed by chromatography using non-phosphopeptide.
Conjugation:	Unconjugated
Storage:	Store the antibody (in aliquots) at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	amyloid beta precursor protein
Database Link:	Entrez Gene 351 Human P05067



[View online »](#)

Background:

Amyloid beta precursor protein gene (ABPP) encodes a cell surface receptor and transmembrane precursor protein that is cleaved by secretases to form a number of peptides. Multiple transcript variants encoding several different isoforms have been found for this gene. Isoform APP695 is the predominant form in neuronal tissue, isoform APP751 and isoform APP770 are widely expressed in nonneuronal cells. Isoform APP751 is the most abundant form in T lymphocytes. ABPP is expressed in all fetal tissues examined with the highest levels in brain, kidney, heart and spleen with weak expression observed in liver; ABPP is induced during neuronal differentiation. In the adult brain, highest expression of ABPP gene is found in the frontal lobe of the cortex and in the anterior perisylvian cortex opercular gyri; moderate expression in the cerebellar cortex, the posterior perisylvian cortex opercular gyri and the temporal associated cortex. Weak expression is found in the striate, extra striate and motor cortices. Mutations in ABPP have been implicated in autosomal dominant Alzheimer disease and cerebroarterial amyloidosis (cerebral amyloid angiopathy).

Synonyms:

Alzheimer disease amyloid protein, Amyloid Precursor Protein, ABPP, APPI, PreA4, Cerebral vascular amyloid peptide, CVAP

Note:

Molecular Weight: 100-140 kDa


Product images:


Figure 1. Western blot analysis of extract from mouse brain tissue, using APP antibody (Lane 1 and 2) and APP (Phospho-Thr668) antibody (Lane 3 and 4).