

Product datasheet for AP02470PU-S

Ezrin (EZR) pThr566 Rabbit Polyclonal Antibody

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: Western Blot: 1:500~1:1000.

Reactivity: Human, Mouse

Host: Rabbit
Clonality: Polyclonal

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human Ezrin

around the phosphorylation site of threonine 566 (Y-K-TP-L-R).

Specificity: Ezrin antibody detects endogenous levels of Ezrin only when phosphorylated at threonine

566.

Formulation: PBS ((without Mg2+ and Ca2+), pH 7.4 containing 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

State: Aff - Purified State: Liquid purified IgG

Concentration: lot specific

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by

chromatography using non-phosphopeptide corresponding to the phosphorylation site.

Conjugation: Unconjugated

Storage: Store the antibody at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: ezrin

Database Link: Entrez Gene 7430 Human

P15311






Background:

Ezrin (also known as p81, 80K, or cytovillin) is a cytoplasmic protein which is enriched in microvilli and other cell surface structures. Its NH2-terminal domain is associated with the plasma membrane and the COOH-terminal domain with the cytoskeleton. Ezrin has an actin-binding capability. Ezrin is expressed in epithelial cells but not in mesenchymal cells. Ezrin is also expressed by certain epithelial tumors, such as renal cell adenocarcinomas. Ezrin serves as a tyrosine kinase substrate, and is phosphorylated in EGF-stimulated cells.

Synonyms:

p81, Cytovillin, Villin-2, EZR, VIL2

Product images:



Western blot analysis of extracts from A431 cells, untreated or EGF-treated (200ng/ml, 30min) using Ezrin antibody (-N, Lane 3 and 4).