

Product datasheet for AP02614PU-S

EGFR Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western Blot: 1/500~1/1000.

Immunohistochemistry: 1/50~1/100.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from human

EGFR around the phosphorylation site of tyrosine 1092 (P-E-YP-I-N).

Specificity: This antibody detects endogenous levels of total EGFR protein.

Formulation: PBS (without Mg2+ and Ca2+), pH 7.4 containing 150mM NaCl, 0.02% Sodium Aazide and 50%

Glycerol.

State: Aff - Purified

State: Liquid purified IgG fraction.

Concentration: lot specific

Purification: Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Storage: Store the antibody (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: epidermal growth factor receptor

Database Link: Entrez Gene 1956 Human

P00533



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. The protein kinase family is one of the largest families of proteins in eukaryotes, classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. Epidermal Growth factor receptor (EGFR) is the prototype member of the type 1 receptor tyrosine kinases. EGFR overexpression in tumors indicates poor prognosis and is observed in tumors of the head and neck, brain, bladder, stomach, breast, lung, endometrium, cervix, vulva, ovary, esophagus, stomach and in squamous cell carcinoma.


Synonyms:

Epidermal growth factor receptor, EGF Receptor, erbB-1, c-ErbB-1

Product images:



Western blot analysis of extracts from HUVEC cells using EGFR antibody (Lane 1 and 2) and EGFR pTyr1092 antibody (Lane 3 and 4).



Immunohistochemical analysis of paraffinembedded human breast carcinoma tissue using EGFR antibody.