

Product datasheet for AP02642PU-S

CDK2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: Western blot (1:500 - 1:1000).

Human, Mouse, Rat Reactivity:

Rabbit Host:

Clonality: Polyclonal

Synthesized non-phosphopeptide derived from human CDK2 around the phosphorylation Immunogen:

site of threonine 160.

Specificity: This antibody detects endogenous levels of total CDK2 protein.

Phosphate buffered saline (without Mg2+and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium Formulation:

> azide and 50% glycerol. State: Aff - Purified

> State: Liquid Ig fraction

Concentration: lot specific

Purification: Affinity-chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store (in aliquots) at -20 °C. Avoid repeated freezing and thawing.

Shelf life: one year from despatch. Stability:

Gene Name: cyclin-dependent kinase 2 Database Link: Entrez Gene 1017 Human

P24941



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Cdk2 is a member of the Ser/Thr protein kinase family. It is highly similar to the gene products of S. cerevisiae cdc28, and S. pombe cdc2. It is a catalytic subunit of the cyclin-dependent protein kinase complex, whose activity is restricted to the G1-S phase, and is essential for cell cycle G1/S phase transition. This protein associates with and is regulated by the regulatory subunits of the complex including cyclin A or E, CDK inhibitor p21Cip1 (CDKN1A) and p27Kip1 (CDKN1B). Its activity is also regulated by protein phosphorylation. Two alternatively spliced variants and multiple transcription initiation sites of this gene have been reported.

Synonyms:

p33 protein kinase

Product images:



Western blot analysis of the extracts from A2780 cells using CDK2 antibody.