

# **Product datasheet for AP02647PU-N**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **MEK3 (MAP2K3) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000. Incubate membrane with diluted antibody in 5% nonfat milk, 1X

TBS, 0,1% Tween-20 at 4°C with gentle shaking, overnight. **Immunohistochemistry on Paraffin Sections:** 1/50-1/100.

Immunofluorescence: 1/100-1/200.

Reactivity: Human, Rat, Mouse

Host: Rabbit
Clonality: Polyclonal

Immunogen: Peptide sequence around amino acids 187~191 (V-D-S-V-A) derived from Human MKK3.

**Specificity:** This MKK3 antibody Cat.-No AP02647PU detects endogenous levels of total MKK3 protein.

Formulation: PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol

State: Aff - Purified

State: Liquid purified Ig fraction

**Concentration:** lot specific

**Purification:** Affinity Chromatography using epitope-specific peptide

Conjugation: Unconjugated

Storage: Upon receipt, store undiluted (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 40 kDa (Predicted)

**Gene Name:** mitogen-activated protein kinase kinase 3

**Database Link:** Entrez Gene 5606 Human

P46734






#### Background:

MEK3 is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is activated by mitogenic and environmental stress, and participates in the MAP kinase-mediated signaling cascade. It phosphorylates and thus activates MAPK14/p38-MAPK. This kinase can be activated by insulin, and is necessary for the expression of glucose transporter. Expression of RAS oncogene is found to result in the accumulation of the active form of this kinase, which thus leads to the constitutive activation of MAPK14, and confers oncogenic transformation of primary cells. The inhibition of this kinase is involved in the pathogenesis of Yersina pseudotuberculosis. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Synonyms:

MAPKK 3, MEK3, MKK3, MAP kinase kinase 3

### **Product images:**



Western Blot analysis of extracts from MDA cells using MKK3 antibody





Figure 1. Immunohistochemical analysis of paraffin- embedded human breast carcinoma tissue, using MKK3 antibody.

Peptide





Immunofluorescence staining of methanol-fixed HeLa cells using MKK3 antibody