

Product datasheet for AP02674PU-N

p27 KIP 1 (CDKN1B) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry: 1/50-1/100.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from human

p27Kip1 around the phosphorylation site of serine 10 (N-G-Sp-P-S).

Specificity: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

This antibody detects endogenous levels of total p27Kip1 protein.

Formulation: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide as preservative and

50% Glycerol as stabilizer.

State: Aff - Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Immunoaffinity Chromatography.

Conjugation: Unconjugated

Storage: Store the antibody (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: cyclin-dependent kinase inhibitor 1B

Database Link: Entrez Gene 1027 Human

P46527



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

p27 is a cyclin dependent kinase inhibitor (CKI) which controls G1/S phase transition in the cell cycle. p27 prevents cell cycle progression by inhibiting the activity of cyclin E-Cdk2 and cyclin D-Cdk4/6, thus preventing E2F activation and S phase progression.

Low p27 levels and/or cytosolic mis-localisation have been associated with the development of lung, breast, colon, oesophagus, and prostate tumours.

Synonyms: p27Kip1

Product images:

Peptide




Figure 2. Western blot analysis using p27Kip1 antibody: Lane 1: The extract from A2780 cells; Lane 2: The extract from COLO205 cells; Lane 3: The extract from COLO205 cells using preincubated with synthesized peptide.




Figure 1. Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using p27Kip1 antibody.