

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for AP02719PU-S

### PYK2 (PTK2B) Rabbit Polyclonal Antibody

### **Product data:**


Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunofluorescence: 1/100-1/200. Immunohistochemistry on Paraffin-Embedded Sections: 1/50-1/100.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The antiserum was produced against synthesized non-phosphopeptide derived from human Pyk2 around the phosphorylation site of tyrosine 402 (D-I-Y <i>P</i> -A-E).
Specificity:	This antibody AP02719PU detects endogenous levels of total Pyk2 protein.
Formulation:	PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol. State: Aff - Purified State: Liquid purified lg fraction.
Concentration:	lot specific
Purification:	Immunoaffinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store the antibody (in aliquots) at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	protein tyrosine kinase 2 beta
Database Link:	<u>Entrez Gene 2185 Human</u> <u>Q14289</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PYK2 (PTK2B) Rabbit Polyclonal Antibody – AP02719PU-S
Background:	PYK2 is involved in calcium induced regulation of ion channel and activation of the map kinase signaling pathway. PKY2 may represent an important signaling intermediate between neuropeptide activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. Interacts with the SH2 domain of Grb2. May phosphorylate the voltage-gated potassium channel protein Kv1.2. PYK2 activation is highly correlated with the stimulation of c-Jun N-terminal kinase activity.
Synonyms:	PYK2, RAFTK, Protein tyrosine kinase 2 beta, Focal adhesion kinase 2, CAK-beta, CADTK


## **Product images:**



Western Blot analysis of extracts from 293 cells using Pyk2 antibody

Immunohistochemical analysis of paraffinembedded human breast carcinoma tissue, using

Pyk2 antibody.




Peptide

.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Immunofluorescence staining of methanol-fixed HeLa cells using Pyk2 antibody (Red).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US