

Product datasheet for **AP03035PU-N**

Acetylated Lysine Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IF, IP, WB
Recommended Dilution:	ELISA. Western blot: A 1/250 dilution of this purified antibody was sufficient to detect the acetylated histone from TSA treated mouse spleen cell in western blot analysis (Ref.6). Immunoprecipitation (Ref.7). Immunofluorescence.
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Acetylated KLH Conjugated
Specificity:	This antibody detects proteins containing Acetylated Lysine residues in SDS-PAGE immunoblots. No reaction to Non-Acetylated proteins.
Formulation:	PBS State: Aff - Purified State: Liquid purified Ig fraction Stabilizer: 50% Glycerol Preservative: 0.09% Sodium Azide
Concentration:	lot specific
Purification:	Affinity Chromatography
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.



[View online »](#)

Background:

Post-translational modifications of proteins play critical roles in the regulation and function of many known biological processes. Proteins can be post-translationally modified in many different ways, and a common posttranscriptional modification of Lysine involves acetylation (1).


The conserved amino-terminal domains of the four core histones (H2A, H2B, H3 and H4) contain lysines that are acetylated by histone acetyltransferases (HATs) and deacetylated by histone deacetylases (HDACs) (2).

Protein posttranslational reversible lysine N ϵ -acetylation and deacetylation have been recognized as an emerging intracellular signaling mechanism that plays critical roles in regulating gene transcription, cell-cycle progression, apoptosis, DNA repair, and cytoskeletal organization (3).


The regulation of protein acetylation status is impaired in the pathologies of cancer and polyglutamine diseases (4), and HDACs have become promising targets for anticancer drugs currently in development (5).

Synonyms:

AcK, acetyl Lysine, acetyl-Lysine

Product images:

Western blot analysis of the acetylated histone from TSA-treated mouse spleen cells.



Western blot analysis of Acetylated Lysine in BSA (Left) and Acetylated BSA (Right) using a 1/1000 dilution of AP03035PU-N