

## Product datasheet for **AP05258PU-N**

### FHL2 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western blot: 0.5-2 µg/ml. Positive control: HeLa cell lysates.
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide derived from the human FHL2 protein
Specificity:	This antibody detects FHL2 / SLIM3. Also binds ABB73038, FHL2 isoform 5 with MW of 389 a.a. (~42kDa).
Formulation:	Phosphate buffered saline with 0.08 % sodium azide State: Aff - Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Ship on dry ice. Upon arrival, freeze in aliquots at -20 °C to -70 °C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	four and a half LIM domains 2
Database Link:	<a href="#">Entrez Gene 2274 Human Q14192</a>

[View online »](#)

**Background:**

Sphingosine kinase-1 (SK1) is a key enzyme that catalyzes the phosphorylation of sphingosine to sphingosine-1-phosphate (S1P). SK1, and its product S1P, regulate numerous and diverse biological functions, including cell growth, differentiation, proliferation, and apoptosis. S1P also plays a central role in cardiac development and ischemic preconditioning. FHL2 (SLIM3) has been identified as a SK1-interacting protein in mammalian cardiomyocytes. FHL2, but not FHL1 or FHL3, interacts with SK1, and FHL2 colocalizes with SK1 in the cytoplasm of these cells. The interaction of FHL2 with SK-1 involves the C-terminal portion of SK1.

Overexpression of FHL2 has been found to attenuate the activity and antiapoptotic effects of SK1. Endothelin-1, a potent survival factor in cardiomyocytes, inhibits FHL2-SK1 association and increases SK1 activity. FHL2 is therefore a novel inhibitor of SK1 activity in cardiomyocytes and targeting FHL2 may prevent myocardial apoptosis through activation of SK1.

A transcriptional coactivator, FHL2 is also known to form complexes consisting of Proline-, glutamic acid-, and leucine-rich protein-1 (PELP1), FHL2, and androgen receptors (AR) in prostate cancer cells, perhaps functioning as a molecular adaptor with PELP1 in the progression of prostate cancer.

**Synonyms:**

FHL-2, SLIM 3, DRAL

**Product images:**


Western blot using, rabbit polyclonal at 0.5 ug/ml on HeLa cell extract (10 ug/lane). Blots were developed with goat anti-rabbit Ig (1:75k) and Pierce's Supersignal West Femto system.