

Product datasheet for AP05264PU-N

CSAG3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: Western blot: 0.5-2 µg/ml.

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide derived from the human Trag-3 protein

Specificity: This antibody detects Trag 3. Trag 3 is weakly expressed in kidney. Expressed in various

tumor cell lines including carcinomas, myeloid and lymphoid malignancies, melanomas and prostate cancer. Overexpressed in taxol-resistant breast cancer line MDA 435(TR) and the

doxorubicin-resistant multiple myelanoma lines 8226/Dox(40) and 8226/MDR(10).

Formulation: Phosphate buffered saline with 0.08 % sodium azide

State: Aff - Purified State: Liquid Ig fraction

Concentration: lot specific

Purification: Affinity chromatography

Conjugation: Unconjugated

Storage: Store the product (in aliquots) at -20 °C. Can be shipped at 2 - 8 °C.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: CSAG family member 3

Database Link: Entrez Gene 389903 Human

Q9Y5P2



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The expression of the cancer-testis antigen Taxol resistance-associated gene-3 (TRAG-3) is associated with acquired paclitaxel (Taxol) resistance, and is expressed in a variety of cancers - e.g., breast cancer, leukemia, and melanoma. Overexpressed in taxol-resistant breast cancer line MDA 435(TR) and the doxorubicin-resistant multiple myelanoma lines 8226/Dox(40) and 8226/MDR(10)V. Weakly expressed in kidney. TRAG-3 is an attractive target for immunotherapy of cancer.

First identified as a novel cancer/testis antigen, TRAG-3, (Taxol Resistance Associated Gene-3) was initially discovered in a search for new genes involved in drug resistance. Early studies of TRAG-3 revealed a minimal to absent expression in normal tissues and a marked over-expression in many carcinoma cell lines including several melanoma lines. By RT-PCR evaluation of TRAG-3 two transcripts are seen in many carcinoma cell lines with products in the the 799 bp and a second alternatively spliced transcript.

Synonyms: TRAG3, CT24.2, TRAG-3

Product images:



Western blot of endogenous Trag3 present in Jurkat cell lysate (15 ug/lane) using (0.5 ug/ml) and developed using anti-rabbit HRP (1:75k) and Pierce's Super Signal West Femto.