

## Product datasheet for AP06010PU-M

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Androgen Receptor (AR) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, IHC

Recommended Dilution: Immunohistochemistry on Paraffin sections: 1/50-1/200.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids 171-220 of Human AR.

**Specificity:** This antibody detects endogenous levels of Androgen receptor protein (region surrounding

Gly207).

**Formulation:** Phosphate buffered saline (PBS), pH 7.2

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% Sodium azide

Concentration: 1.0 mg/ml

**Purification:** Affinity chromatography using epitope-specific immunogen (> 95% pure; by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 110 kDa

**Gene Name:** androgen receptor

Database Link: Entrez Gene 367 Human

P10275






Background:

Androgens exhibit a wide range of effects on the development, maintenance and regulation of male phenotype and make reproductive physiology. The androgen receptor (AR) is a member of the steroid superfamily of liganddependent transcription factors. ARs bind the two biologically active androgens, testosterone (T) and dihydrotestosterone (DHT), with high and nearly identical affinities; however, the rates of association and dissociation of T are about three times more rapid than those of DHT. This difference has resulted in speculation as to whether these differences in binding kinetics could account for the different physiological effects of T and DHT. A striking feature of AR is its rapid degradation in the absence of ligand. It is now well established that androgen binding results in an at least sixfold increase in androgen stability and that ligand-induced stabilization of AR is highly androgen-specific.

Synonyms:

Dihydrotestosterone receptor, DHTR, NR3C4

## **Product images:**



Immunohistochemical analysis using Androgen receptor antibody in Paraffin-embedded human prostate carcinoma tissue.