

Product datasheet for AP06010PU-N

Androgen Receptor (AR) Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	ELISA, IHC
Recommended Dilution:	Immunohistochemistry on Paraffin sections 1/50-1/200.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 171-220 of Human AR.
Specificity:	This antibody detects endogenous levels of Androgen receptor protein (region surrounding Gly207).
Formulation:	Phosphate buffered saline (PBS), pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% Sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity chromatography using epitope-specific immunogen (> 95% pure; by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 110 kDa
Gene Name:	androgen receptor
Database Link:	<u>Entrez Gene 367 Human</u> <u>P10275</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Serigene Androgen Receptor (AR) Rabbit Polyclonal Antibody – AP06010PU-N

Background: Androgens exhibit a wide range of effects on the development, maintenance and regulation of male phenotype and make reproductive physiology. The androgen receptor (AR) is a member of the steroid superfamily of liganddependent transcription factors. ARs bind the two biologically active androgens, testosterone (T) and dihydrotestosterone (DHT), with high and nearly identical affinities; however, the rates of association and dissociation of T are about three times more rapid than those of DHT. This difference has resulted in speculation as to whether these differences in binding kinetics could account for the different physiological effects of T and DHT. A striking feature of AR is its rapid degradation in the absence of ligand. It is now well established that androgen binding results in an at least sixfold increase in androgen stability and that ligand-induced stabilization of AR is highly androgen- specific.

Synonyms:

Dihydrotestosterone receptor, DHTR, NR3C4

Product images:



Immunohistochemical analysis using Androgen receptor antibody in Paraffin-embedded human prostate carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US