

# **Product datasheet for AP06022PU-N**

# Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

## **ATF2 Rabbit Polyclonal Antibody**

**Product data:** 

Product Type: Primary Antibodies

Applications: ELISA, IHC, IP, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on Paraffin sections: 1/50-1/200.

**Immunoprecipitation**: 1/50-1/200.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids 30-80 of Human ATF2.

**Specificity:** This antibody detects endogenous levels of ATF2 protein (region surrounding Val65/47).

**Formulation:** Phosphate buffered saline (PBS), pH 7.2

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% Sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity chromatography using epitope-specific immunogen (> 95% pure by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~55 to 75 kDa

**Gene Name:** activating transcription factor 2

**Database Link:** Entrez Gene 1386 Human

P15336






### Background:

The transcription factor ATF-2 (also called CRE-BP1) binds to both AP-1 and CRE DNA response elements and is a member of the ATF/CREB family of leucine zipper proteins. ATF-2 interacts with a variety of viral oncoproteins and cellular tumor suppressors and is a target of the SAPK/JNK and p38 MAP kinase signaling pathways. Various forms of cellular stress, including genotoxic agents, inflammatory cytokines and UV irradiation, stimulate the transcriptional activity of ATF-2. Cellular stress activates ATF-2 by phosphorylation of Thr69 and Thr71. Both SAPK and p38 MAPK have been shown to phosphorylate ATF-2 at these sites in vitro and in cells transfected with ATF-2.


Synonyms:

ATF-2, CREB2, CREBP1, Activating transcription factor 2, CRE-BP1, HB16, CREB-2

### **Product images:**



Western blot (WB) analysis of ATF2 antibody at 1/500 dilution Lane 1:Hela cell lysate Lane 2:Mouse kidney tissue lysate Lane 3:Rat brain tissue lysate



Immunohistochemical analysis in paraffinembedded human breast carcinoma tissue using ATF2 antibody.