

Product datasheet for AP06032PU-N

Bim (BCL2L11) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

ELISA, IHC, WB **Applications:**

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on Paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Synthetic peptide, corresponding to amino acids 1-50 of Human Bim. Immunogen:

This antibody detects endogenous levels of Bcl-2-like 11 / BIM-α/β protein (region Specificity:

surrounding Ala2).

Formulation: Phosphate buffered saline (PBS), pH 7.2

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% Sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity chromatography using epitope-specific immunogen (> 95% pure by SDS-PAGE).

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~25 kDa

Gene Name: BCL2 like 11

Database Link: Entrez Gene 10018 Human

O43521



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Pro-apototic Bcl-2 family members promote cell death by neutralizing their anti-apoptotic relatives, which otherwise maintain cell viability by regulating caspase activity. Bim belongs to the BH3-only subgroup of Bcl-2 related proteins and exists in three distinct isoforms, BimS (short), BimL (long) and BimEL (extra long). ERK1/2 phosphorylates BimEL, resulting in rapid degradation of the isoform via the proteasome pathway. At least three sites for ERK1/2 phosphorylation exist on BimEL, whereas ERK1/2 does not effect BimS or BimL, implying a unique role for BimEL in cell survival signaling.


Synonyms:

Bcl2-L-11, BCL2L11, BIM, BimEL, BimL, BimS

Product images:



Western blot (WB) analysis of Bim antibody at 1/500 dilution Lane 1:Hela cell lysate Lane 2:Mouse kidney tissue lysate Lane 3:Rat kidney tissue lysate



Immunohistochemical analysis in paraffinembedded human breast carcinoma tissue using Bcl-2-like 11 antibody.