

# Product datasheet for AP06038PU-M

## beta Catenin (CTNNB1) Rabbit Polyclonal Antibody

### **Product data:**

#### **Product Type: Primary Antibodies** ELISA, IHC, WB **Applications:** Recommended Dilution: Western blot: 1/500-1/1000. Human, Mouse, Rat **Reactivity:** Host: Rabbit **Clonality:** Polyclonal Immunogen: Synthetic peptide, corresponding to amino acids N-terminus of Human Catenin-β. Specificity: This antibody detects endogenous levels of Catenin- $\beta$ protein (region surrounding lle35). Formulation: Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified lg fraction Preservative: 0.05% sodium azide Concentration: 1.0 mg/ml **Purification:** Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE) **Conjugation:** Unconjugated Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Stability: Shelf life: one year from despatch. **Predicted Protein Size:** ~ 92 kDa Gene Name: catenin beta 1 Database Link: Entrez Gene 1499 Human P35222



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


### Seta Catenin (CTNNB1) Rabbit Polyclonal Antibody – AP06038PU-M

Background:The catenins, α, β and γ, are proteins which bind to the highly conserved, intracellular<br/>cytoplasmic tail of E-cadherin. Together, the catenin/cadherin complexes play an important<br/>role mediating cellular adhesion. α-catenin was initially described as an E-cadherin associated<br/>protein, and since has been shown to associate with other members of the cadherin family,<br/>such as N-cadherin and P-cadherin. β-catenin associates with the cytoplasmic portion of E-<br/>cadherin, which is necessary for the function of E-cadherin as an adhesion molecule. β-<br/>catenin has also been found in complexes with the tumor suppressor protein APC. γ-catenin,<br/>also known as plakoglobin, is a protein that binds with α-catenin and N-cadherin. It has been<br/>shown thatthe transmembrane phosphatase PTPµ associates with catenin/cadherin<br/>complexes and may regulate complex signaling.


Synonyms:

CTNNB1, CTNNB, Beta-catenin

### **Product images:**



Western blot analysis in extracts from SW626 cell using Catenin- $\beta$  antibody.



Immunohistochemistry (IHC) analysis of Catenin- $\beta$  antibody on paraffin-embedded sections.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US