

Product datasheet for AP06051PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

Chk2 (CHEK2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

WB **Applications:**

Western blot: 1/500-1/1000. **Recommended Dilution:**

Reactivity: Human

Rabbit Host:

Clonality: Polyclonal

Synthetic peptide, corresponding to amino acids 471-520 of Human Chk2. Immunogen:

This antibody detects endogenous levels of Chk2 protein (region surrounding Gln510). **Specificity:**

Phosphate buffered saline (PBS), pH 7.2. Formulation:

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

Concentration: 1.0 ma/ml

Purification: Affinity chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Unconjugated Conjugation:

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Shelf life: one year from despatch. Stability:

~ 62 kDa **Predicted Protein Size:**

checkpoint kinase 2 Gene Name:

Database Link: Entrez Gene 11200 Human

096017






Background:

Cell cycle events are regulated by the sequential activation and deactivation of cyclin dependent kinases (Cdks) and by proteolysis of cyclins. Chkl and Chk2 are involved in these processes as regulators of Cdks. Chkl and Chk2 both function as essential components in the G2 DNA damage checkpoint by phosphorylating Cdc25C in response to DNA damage. Phosphorylation inhibits Cdc25C activity, thereby blocking mitosis. Cdc25A, Cdc25B and Cdc25C protein tyrosine phosphatases function as mitotic activators by dephosphorylating Cdc2 p34 on regulatory tyrosine residues. It has also been shown that Chkl can phosphorylate Weel in vitro, providing evidence that the hyperphosphorylated form of Weel, seen in cells delayed by Chkl overexpression, is due to phosphorylation by Chkl.

Synonyms:

CHEK2, CHEK-2, CHK-2, RAD53, Cds1

Product images:



Western blot (WB) analysis of Chk2 antibody at 1/500 dilution Lane 1:Hela cell lysate Lane 2:Mouse liver tissue lysate Lane 3:Rat spleen tissue lysate