

Product datasheet for AP06056PU-S

c-Jun (JUN) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

IHC, WB **Applications:**

Recommended Dilution: Immunohistochemistry on Paraffin sections 1/50-1/200

Western blot: 1/500-1/1000.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Synthetic peptide, corresponding to amino acids 201-250 of Human c-Jun. Immunogen:

Antibody detects endogenous levels of AP-1 / c-lun protein (region surrounding Asn57). Specificity:

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% Sodium azide

Concentration: 1.0 mg/ml

Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-**Purification:**

PAGE)

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 43, 48 kDa

Gene Name: Jun proto-oncogene, AP-1 transcription factor subunit

Database Link: Entrez Gene 3725 Human

P05412



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Genes belonging to the Jun and Fos oncogene families encode nuclear proteins that are found to be associated with a number of transcriptional complexes. The c-Jun protein is a major component of the transcription factor AP-1, originally shown to mediate phorbol ester tumor promoter (TPA)-induced expression of responsive genes through the TPA- response element (TRE). The Jun proteins form homo- and heterodimers which bind the TRE, while Fos proteins are active only as heterodimers with any of the Jun proteins. Fos/Jun heterodimers have a much higher affinity for the TRE than Jun homodimers. Ha-Ras augments c-Jun activity and stimulates phosphorylation of its activation domain. An inhibitor of Fos/Jun function, termed IP-1, associates with Fos and Jun and is inactivated upon phosphorylation induced by the cAMP-dependent protein kinase A (PKA).

Synonyms:

Transcription factor AP1

Product images:



Western blot (WB) analysis of c-Jun antibody at 1/500 dilution Lane 1:HEK293T cell lysate Lane 2:Raw264.7 cell lysate Lane 3:Rat kidney tissue lysate