

Product datasheet for AP06086PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

PARK7 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on Paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of DJ-1/PARK7 protein (region surrounding Pro54).

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% Sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity chromatography using epitope-specific immunogen (> 95% pure by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~22 kDa

Gene Name: Parkinsonism associated deglycase

Database Link: Entrez Gene 11315 Human

Q99497






Background:

The DJ-1 gene encodes a highly-conserved protein which is implicated in a number of cell processes. DJ-1 was first identified as a novel oncogene that transformed mouse NIH/3T3 cells in cooperation with activated Ras. Additionally, DJ-1 was cloned in rat as SP22 or CAP-1 and found to be an infertilityrelated sperm protein, whose expression is reduced in sperm treated with toxicants. DJ-1 also positively regulates the androgen receptor (AR) by forming a complex with PIASx α , a negative regulator of AR. The gene encoding human DJ-1 maps to chromosome 1p36.33-p36.12, a region identified as a hot spot of chromosome abnormalities in several tumor cells. Subsequently, mutations in the DJ-1 gene have been implicated in Parkinson's disease, and loss of DJ-1 function leads to neurodegeneration. DJ-1 is an ubiquitously expressed protein that is induced in response to growth stimuli and translocates from the cytoplasm to the nucleus during the S phase of the cell cycle.


Synonyms:

Oncogene DJ1, Parkinson disease protein 7

Product images:



Western blot analysis in extracts from HuvEc cells using DJ-1 / PARK7 antibody (Cat.-No. AP06086PU-N).



Immunohistochemical analysis in paraffinembedded human lung carcinoma tissue using DJ-1 / PARK7 antibody (Cat.-No. AP06086PU-N).