

Product datasheet for AP06087PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DNA Polymerase beta (POLB) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on Paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

Specificity: This antibody detects endogenous levels of DNA polymerase beta / DPOLB protein (region

surrounding Gln324).

Formulation: Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~38 kDa

Gene Name: polymerase (DNA) beta

Database Link: Entrez Gene 18970 MouseEntrez Gene 29240 RatEntrez Gene 5423 Human

P06746




Background:

DNA replication, recombination and repair, all of which are necessary for genomic stability, require the presence of exonucleases. In DNA replication, these enzymes are involved in the processing of Okazaki fragments, whereas in DNA repair they function to excise damaged DNA fragments and correct recombinational mismatches. These exonucleases include the family of DNA polymerases. DNA pol α , β , δ and ϵ are involved in DNA replication and repair. DNA pol δ and DNA pol ϵ are multisubunit enzymes, with DNA pol δ consisting of two subunits—p125, which interacts with the sliding DNA clamp protein PCNA, and p50. The nuclear-encoded DNA pol γ is the only DNA polymerase required for the replication of the mitochondrial DNA. DNA pol ζ is ubiquitously expressed in various tissues and mediates the cellular mechanism of damage-induced mutagenesis. DNA pol θ is a DNA polymerase-helicase that binds ATP and is involved in the repair of interstrand crosslinks.

Synonyms: POLB


Protein Families: Druggable Genome
Protein Pathways: Base excision repair

Product images:



Western blot (WB) analysis of DNA pol β antibody (Cat.-No.: [AP06087PU-N]) at 1/500 dilution Lane 1:Hela cell lysate Lane 2:sp2/0 cell lysate Lane 3:Rat brain tissue lysate





Immunohistochemical analysis in paraffinembedded human breast carcinoma tissue using DNA polymerase beta Antibody Cat.-No [AP06087PU-N].